

1 STATE-SPACE MODELLING OF DATA ON MARKED  
2 INDIVIDUALS

3 Olivier GIMENEZ<sup>1,2,6</sup>, Vivien ROSSI<sup>3,4,5</sup>, Rémi CHOQUET<sup>1</sup>, Camille DEHAIS<sup>1</sup>,  
4 Blaise DORIS<sup>1,3</sup>, Hubert VARELLA<sup>1,3</sup>, Jean-Pierre VILA<sup>3</sup> and Roger PRADEL<sup>1</sup>

5 <sup>1</sup>*Centre d'Ecologie Fonctionnelle et Evolutive/CNRS - UMR 5175*

6 *1919 Route de Mende*

7 *34293 Montpellier - FRANCE*

8 <sup>2</sup>*Centre for Research into Ecological and Environmental Modelling*

9 *University of St Andrews, St Andrews*

10 *The Observatory, Buchanan Gardens, KY16 9LZ - SCOTLAND*

11 <sup>3</sup>*Laboratoire d'Analyse des Systèmes et Biométrie - UMR 729*

12 *INRA/ENSAM*

13 *2 Place Pierre Viala, 34060 Montpellier - FRANCE*

14 <sup>4</sup>*Institut de Modélisation Mathématiques de Montpellier - UMR 5149*

15 *Universite Montpellier 2*

16 *CC051, Place Eugène Bataillon, 34095 Montpellier Cedex 5 - FRANCE*

17 <sup>5</sup>*CIRAD - Unité de dynamique des forêts naturelles*

18 *TA C-37 / D*

19 *Campus International de Baillarguet, 3434398 Montpellier Cedex 5 - FRANCE*

20 <sup>6</sup>*E-mail: olivier.gimenez@cefe.cnrs.fr*

21 **Abstract**

22 State-space models have recently been proposed as a convenient and flexible  
23 framework for specifying stochastic models for the dynamics of wild animal pop-  
24 ulations. Here we focus on the modelling of data on marked individuals which

1 is frequently used in order to estimate demographic parameters while accounting  
2 for imperfect detectability. We show how usual models to deal with capture-  
3 recapture and ring-recovery data can be fruitfully written as state-space models.  
4 An illustration is given using real data and a Bayesian approach using MCMC  
5 methods is implemented to estimate the parameters. Eventually, we discuss fu-  
6 ture developments that may be facilitated by the SSM formulation.

## 7 **1 Introduction**

8 The estimation of animal survival is essential in population biology to investigate pop-  
9 ulation dynamics, with important applications in the understanding of ecological, evo-  
10 lutionary, conservation and management issues for wild populations (Pollock, 1991;  
11 Williams et al., 2002). While the time to event is known in medical, social or engi-  
12 neering sciences (death, marriage and failure respectively), models for estimating wild  
13 animal survival must incorporate nuisance parameters to account for incomplete de-  
14 tectability in monitoring individuals (Schwarz et Seber, 1999). Typically, individuals  
15 are captured, marked and can be resighted or recaptured (encountered thereafter) to  
16 construct encounter histories which consist of sequences of 1's and 0's according to  
17 whether a detection occurs or not. The likelihood for such data arises from products  
18 of multinomial distributions whose cell probabilities are complex functions of survival  
19 probabilities - parameters of primary interest - and encounter probabilities - nuisance  
20 parameters (Cormack, 1964; Jolly, 1965; Seber, 1965 - CJS thereafter).

21 In this note, we show how the population process can be fruitfully disentangled,  
22 by distinguishing the underlying demographic process, i.e. the survival (as well as  
23 transitions between sites/states if needed), from its observation, i.e. the detectability.  
24 This leads us to consider a natural formulation for capture-recapture models using state-

1 space models (SSMs). Our contribution is in line with a recent paper by Buckland et  
2 al. (2004) who have proposed to adopt SSMs as a convenient and flexible framework  
3 for specifying stochastic models for the dynamics of wild animal populations.

4 Thus far, SSMs have been mainly used for time series of animal counts (de Valpine,  
5 2004; Millar and Meyer, 2000) or animal locations (Anderson-Sprecher and Ledolter,  
6 1991) to allow true but unobservable states (the population size or trajectory) to be  
7 inferred from observed but noisy data (see Clark et al., 2005 and Wang, 2006 for  
8 reviews). The novelty of our approach lies in the use of SSMs to fit capture-recapture  
9 models to encounter histories.

10 In Section 2, we discuss how to express the CJS model under the form of a SSM. The  
11 implementation details are provided, and real data are presented to compare parameter  
12 estimates as obtained using the standard product-multinomial and the SSM approaches.  
13 In Section 3, the flexibility of the state-space modeling approach is demonstrated by  
14 considering two widely used alternative schemes for collecting data on marked animals.  
15 Finally, Section 4 discusses important developments of capture-recapture models facil-  
16 itated by the SSM formulation. We emphasize that this general framework has a great  
17 potential in population ecology modelling.

## 18 **2 State-space modelling of capture-recapture data**

19 We focus here on the CJS model for estimating animal survival based on capture-  
20 recapture data, as this model is widely used in the ecological and evolutionary literature  
21 (e.g. Lebreton et al., 1992).

## 1 2.1 Likelihood

2 We first define the observations and then the states of the system. We assume that  
3  $n$  individuals are involved in the study with  $T$  encounter occasions. Let  $X_{i,t}$  be the  
4 binary random variable taking values 1 if individual  $i$  is alive at time  $t$  and 0 if it is  
5 dead at time  $t$ . Let  $Y_{i,t}$  be the binary random variable taking values 1 if individual  $i$  is  
6 encountered at time  $t$  and 0 otherwise. Note that we consider the encounter event as  
7 being physically captured or barely observed. The parameters involved in the likelihood  
8 are  $\phi_{i,t}$ , the probability that an animal  $i$  survives to time  $t + 1$  given that it is alive  
9 at time  $t$  ( $t = 1, \dots, T - 1$ ), and  $p_{i,t}$  the probability of detecting individual  $i$  at time  $t$   
10 ( $t = 2, \dots, T$ ). Let finally  $e_i$  be the occasion where individual  $i$  is encountered for the  
11 first time. A general state-space formulation of the CJS model is therefore given by:

$$12 \quad Y_{i,t}|X_{i,t} \sim \text{Bernoulli}(X_{i,t}p_{i,t}), \quad (1)$$

$$13 \quad X_{i,t+1}|X_{i,t} \sim \text{Bernoulli}(X_{i,t}\phi_{i,t}), \quad (2)$$

14 for  $t \geq e_i$ , with  $p_{i,e_i} = 1$  and where Equation (1) and Equation (2) are the observation  
15 and the state equations respectively. This formulation naturally separates the nuisance  
16 parameters (the encounter probabilities) from the parameters of actual interest i.e. the  
17 survival probabilities, the latter being involved exclusively in the state Equation (2).  
18 Such a clear distinction between a demographic process and its observation makes the  
19 description of a biological dynamic system much simpler and allows complex models  
20 to be fitted (Pradel, 2005; Clark et al., 2005). We will refer to this formulation as  
21 the individual state-space CJS model (individual SSM CJS hereafter). The rationale  
22 behind the above formulation is as follows. We give the full details for the observation  
23 Equation (1) only, as a similar reasoning easily leads to Equation (2). If individual  $i$  is

1 alive at time  $t$ , then it has probability  $p_{i,t}$  of being encountered and probability  $1 - p_{i,t}$   
2 otherwise, which translates into  $Y_{i,t}$  is distributed as  $Bernoulli(p_{i,t})$  given  $X_{i,t} = 1$ . Now  
3 if individual  $i$  is dead at time  $t$ , then it cannot be encountered, which translates into  
4  $Y_{i,t}$  is distributed as  $Bernoulli(0)$  given  $X_{i,t} = 0$ . Putting together those two pieces of  
5 reasoning, the distribution of the observation  $Y_{i,t}$  conditional on the state  $X_{i,t}$  is given  
6 by Equation (1).

7 Statistical inference then requires the likelihood of the state-space model specified  
8 above. Assuming independence of individuals, the likelihood is given by the product  
9 of all individual likelihood components. The likelihood component for individual  $i$  is  
10 the probability of the vector of observations  $\mathbf{Y}_i^T = (Y_{i,e_i}, \dots, Y_{i,T})$  which gathers the  
11 information set up to time  $T$  for this particular individual. Conditional on the first  
12 detection, the likelihood component corresponding to individual  $i$  is therefore given by  
13 (e.g. Harvey, 1989)

$$14 \quad \int_{X_{i,e_i}} \dots \int_{X_{i,T}} [X_{i,e_i}] \left\{ \prod_{t=e_i+1}^T [Y_{i,t}|X_{i,t}][X_{i,t}|X_{i,t-1}] \right\} dX_{i,e_i} \dots dX_{i,T} \quad (3)$$

15 where  $[X]$  denotes the distribution of  $X$  and  $X_{i,e_i}$  the initial state of individual  $i$  which  
16 is assumed to be alive. Because we deal with binary random vectors, we used the  
17 counting measure instead of the Lebesgue measure.

18 In its original formulation, the CJS makes important assumptions regarding indi-  
19 viduals. First, all individuals share the same parameters, which means that the survival  
20 and detection probabilities depend on the time index only. In mathematical notation,  
21 we have  $\phi_{i,t} = \phi_t$  and  $p_{i,t} = p_t$  for all  $i = 1, \dots, n$ , so Equation (1) and Equation (2)  
22 become  $X_{i,t+1}|X_{i,t} \sim Bernoulli(X_{i,t}\phi_t)$  and  $Y_{i,t}|X_{i,t} \sim Bernoulli(X_{i,t}p_t)$  respectively.  
23 Second, the CJS model also assumes independence between individuals. By using sim-

1 ple relationships between Bernoulli and Binomial distributions, one can finally fruitfully  
 2 formulates the original CJS model under the following state-space model:

$$3 \quad Y_t|X_t \sim \text{Bin}(X_t - u_t, p_t) \quad (4)$$

$$4 \quad X_{t+1}|X_t \sim \text{Bin}(X_t, \phi_t) + u_{t+1} \quad (5)$$

5 where  $X_t$  is the number of survivors from time  $t$  plus the number of newly marked  
 6 individuals at time  $t$ ,  $u_t$ , and  $Y_t$  is the total number of previously marked individuals  
 7 encountered at time  $t$ . We will refer to this formulation as the population state-space  
 8 CJS model (population SSM CJS hereafter). Interestingly, specifying the system un-  
 9 der a state-space formulation now requires much less equations than the individual  
 10 SSM CJS model, which may avoid the computational burden. Nevertheless, while the  
 11 individual SSM CJS involves parameters for every single individual and sampling occa-  
 12 sion, the population SSM CJS model makes the strong assumptions that all individuals  
 13 behave the same as well as independently, which may be of little relevance from the  
 14 biological point of view. To cope with this issue, in-between modelling can be achieved  
 15 by considering age effects or groups classes in the population SSM model (Lebreton et  
 16 al., 1992). Finally, covariates can be incorporated in order to assess the effect of envi-  
 17 ronment such as climate change, most conveniently by writing the relationship between  
 18 the target probabilities and the predictors on the logit scale (Pollock, 2002).

## 19 **2.2 Implementation**

20 Fitting SSMs is complicated due to the high-dimensional integral involved in the indi-  
 21 vidual likelihood Equation (3). To circumvent this issue, several techniques have been  
 22 proposed including Kalman filtering, Monte-Carlo particle filtering (such as sequential

1 importance sampling) and MCMC (see Clark et al., 2005 and Wang, 2006 for reviews).  
2 Our objective here is not to discuss these different methods. For our purpose, we adopt  
3 the MCMC technique which is now widely used in biology (Ellison, 2004; Clark, 2005),  
4 in particular for estimating animal survival (Seber et Schwarz, 1999; Brooks et al.,  
5 2000). Besides, this is to our knowledge the only methodology which comes with an  
6 efficient and flexible program to implement it, which, in our case, will allow biologists  
7 to easily and rapidly adopt our approach.

8 In addition to the difficulty of estimating model parameters, the use of SSMS raises  
9 several important issues regarding identifiability, model selection and goodness-of-fit  
10 (Buckland et al. 2004) that were not discussed here. Noteworthy, Bayesian modelling  
11 using MCMC methods offer possible solutions reviewed in Gimenez et al. (submitted).

## 12 **2.3 Illustration**

13 We consider capture-recapture data on the European dipper (*Cinclus cinclus*) that were  
14 collected between 1981 and 1987 (Lebreton et al., 1992). The data consists of marking  
15 and recaptures of 294 birds ringed as adults in eastern France. We applied standard  
16 maximum-likelihood estimation (Lebreton et al. 1992) and MCMC techniques (Brooks  
17 et al. 2000) using the product-multinomial likelihood and the state-space likelihood of  
18 Equation (3) in combination with Equation (1) and Equation (2). We ran two overdis-  
19 persed parallel MCMC chains to check whether convergence was reached (Gelman,  
20 1996). We used 10,000 iterations with 5,000 burned iterations for posterior summariza-  
21 tion. We used uniform flat priors for both survival and detection probabilities. The  
22 simulations were performed using WinBUGS (Spiegelhalter et al., 2003). The R (Ihaka  
23 and Gentleman, 1996) package R2WinBUGS (Sturtz et al., 2005) was used to call Win-  
24 BUGS and export results in R. This was especially helpful when converting the raw

1 encounter histories into the appropriate format, generating initial values and calculate  
2 posterior modes. The programs are available in Appendix A. Posterior summaries for  
3 encounter and survival probabilities are given in Table 1, along with their posterior  
4 probability distributions that are displayed in Figure 1.

5 [Table 1 about here.]

6 [Figure 1 about here.]

7 Survival estimates were uniformly similar whatever the method used (Table 1). In  
8 particular, there is a clear decrease in survival 1982-1983 and 1983-1984, corresponding  
9 to a major flood during the breeding season in 1983 (Lebreton et al., 1992).

10 In contrast, posterior medians of detection probabilities using the CJS SSM ap-  
11 proach are quite different from the classical maximum likelihood estimates, but more  
12 similar to the posterior medians obtained with the product-multinomial likelihood ap-  
13 proach (Table 1). These discrepancies are no longer present when posterior modes  
14 are examined, as expected (recall that we use non-informative uniform distributions as  
15 priors for all parameters).

16 The last survival probability as well as the last detection probability are estimated  
17 with high variability (Table 1 and Figure 1). The fact that these two parameters cannot  
18 be separately estimated is not surprising since the CJS model is known to be parameter-  
19 redundant (Catchpole and Morgan, 1997). Also, the first survival probability and the  
20 first detection probability are poorly estimated, due to the fact that very few individuals  
21 were marked at the first sampling occasion (approximately 7% of the full data set).

22 In terms of time computation, the MCMC approach using a product-multinomial  
23 likelihood took 30 seconds to run and a few second for the classical approach, while



1 the MCMC approach using the SSM likelihood took 4 minutes (512Mo RAM, 2.6GHz  
 2 CPU).

### 3 **3 Further state-space modelling**

#### 4 **3.1 Multistate capture-recapture models**

5 Multistate capture-recapture models (Arnason, 1973; Schwarz et al., 1993; AS hereafter)  
 6 are a natural generalization of the CJS model in that individuals can move between  
 7 states, according to probabilities of transition between those states. States can be either  
 8 geographical sites or states of categorical variables like reproductive status or size class  
 9 (Lebreton and Pradel, 2002). We provide here a state-space modelling formulation  
 10 of the AS model (Dupuis, 1995; Newman, 1998; Clark et al., 2005). Without loss of  
 11 generality, we consider 2 states. Let  $X_{i,t}$  be the random state vector taking values  
 12  $(1, 0, 0)$ ,  $(0, 1, 0)$  and  $(0, 0, 1)$  if, at time  $t$ , individual  $i$  is alive in state 1, 2 or dead  
 13 respectively. Let  $Y_{i,t}$  be the random observation vector taking values  $(1, 0, 0)$ ,  $(0, 1, 0)$   
 14 and  $(0, 0, 1)$  if, at time  $t$ , individual  $i$  is encountered in state 1, 2 or not encountered.  
 15 Parameters involved in the modelling include  $\phi_{i,t}^{rs}$ , the probability that an animal  $i$   
 16 survives to time  $t + 1$  given that it is alive at time  $t$  ( $t = 1, \dots, T - 1$ ) and makes the  
 17 transition between state  $r$  and state  $s$  over the same interval ( $r, s = 1, 2$ ), as well as  $p_{i,t}^r$   
 18 the probability of detecting individual  $i$  at time  $t$  in state  $r$  ( $t = 2, \dots, T$ ,  $r = 1, 2$ ). A  
 19 state-space formulation for the multistate AS model is given by:

$$Y_{i,t}|X_{i,t} \sim \text{Multinomial} \left( 1, X_{i,t} \begin{bmatrix} p_{i,t}^1 & 0 & 1 - p_{i,t}^1 \\ 0 & p_{i,t}^2 & 1 - p_{i,t}^2 \\ 0 & 0 & 1 \end{bmatrix} \right) \quad (6)$$

20

$$X_{i,t+1}|X_{i,t} \sim \text{Multinomial} \left( 1, X_{i,t} \begin{bmatrix} \phi_{i,t}^{11} & \phi_{i,t}^{12} & 1 - \phi_{i,t}^{11} - \phi_{i,t}^{12} \\ \phi_{i,t}^{21} & \phi_{i,t}^{22} & 1 - \phi_{i,t}^{21} - \phi_{i,t}^{22} \\ 0 & 0 & 1 \end{bmatrix} \right) \quad (7)$$

2 where Equation (6) and Equation (7) are the observation and the state equations  
 3 respectively. This formulation has similarities with that of Pradel (2005) who used  
 4 hidden-Markov models to extend multistate models to cope with uncertainty in state  
 5 assignment. Again, it should be noted that the state-space formulation allows the de-  
 6 mographic parameters to be distinguished from the nuisance parameters. A similar  
 7 reasoning to that adopted for the CJS model leads to Equations (6) and (7). As ex-  
 8 pected, Equation (6) and Equation (7) reduce to Equation (1) and Equation (2) if one  
 9 considers a single state. Making similar assumptions as in the CJS model leads to the  
 10 population AS SSM.

## 11 **3.2 Ring-recovery models**

12 The capture-recapture models presented above deals with apparent survival, which  
 13 turns out to be true survival if emigration is negligible. When marks of individuals  
 14 (or individuals themselves) are actually recovered, true survival probabilities can be  
 15 estimated using ring-recovery models (Brownie et al., 1985; RR models hereafter). Let  
 16  $X_{i,t}$  be the binary random variable taking values 1 if individual  $i$  is alive at time  $t$  and  
 17 0 if it is dead at time  $t$ . Let  $Y_{i,t}$  be the binary random variable taking values 1 if mark  
 18 of individual  $i$  is recovered at time  $t$  and 0 otherwise. The parameters involved in the  
 19 likelihood are  $\phi_{i,t}$ , the probability that an animal  $i$  survives to time  $t + 1$  given that  
 20 it is alive at time  $t$  ( $t = 1, \dots, T - 1$ ), and  $\lambda_{i,t}$  the probability of recovering the mark  
 21 of individual  $i$  at time  $t$  ( $t = 2, \dots, T$ ). A general state-space formulation of the RR

1 model is therefore given by:

$$2 \quad Y_{i,t}|X_{i,t}, X_{i,t-1} \sim \text{Bernoulli}((X_{i,t-1} - X_{i,t})\lambda_{i,t}) \quad (8)$$

$$3 \quad X_{i,t+1}|X_{i,t} \sim \text{Bernoulli}(X_{i,t}\phi_{i,t}) \quad (9)$$

4 where Equation (8) and Equation (9) are the observation and the state equations re-  
5 spectively. While the state Equation (9) is the same as that in the individual SSM  
6 CJS, the observation Equation (8) deserves further explanation. If individual  $i$ , alive  
7 at time  $t$ , does not survive to time  $t + 1$ , then its mark has probability  $\lambda_{i,t}$  of being  
8 recovered and probability  $1 - \lambda_{i,t}$  otherwise, which translates into  $Y_{i,t}$  is distributed as  
9  $\text{Bernoulli}(\lambda_{i,t})$  given  $X_{i,t-1} = 1$  and  $X_{i,t} = 0$  i.e.  $X_{i,t-1} - X_{i,t} = 1$ . Now if individual  $i$   
10 is in a given state (dead or alive) at time  $t$  and remains in this state till time  $t + 1$ , then  
11 its mark cannot be recovered, which translates into  $Y_{i,t}$  is distributed as  $\text{Bernoulli}(0)$   
12 given  $X_{i,t-1} = 0$  and  $X_{i,t} = 0$  or  $X_{i,t-1} = 1$  and  $X_{i,t} = 1$  i.e.  $X_{i,t-1} - X_{i,t} = 0$ . The dis-  
13 tribution of the observation  $Y_{i,t}$  conditional on the combination of states  $X_{i,t-1} - X_{i,t}$  is  
14 thus given by Equation (8). Similar comments to that of previous sections can be made  
15 here as well. Finally, we note that because the probability distribution of the current  
16 observation does not only depend on the current state variable, the model defined by  
17 Equation (8) and Equation (9) does not exactly matches the definition of a state-space  
18 model but can be rewritten as such (see Appendix B).

## 19 4 Discussion

20 We have shown that, by separating the demographic process from its observation, CR  
21 models for estimating survival can be expressed as SSMs. In particular, the SSM  
22 formulation of the CJS model competes well with the standard method when applied to

1 a real data set. Bearing this in mind, we see at least two further promising developments  
2 to our approach.

3 First, by scaling down from the population to the individual level while modelling  
4 the survival probabilities, individual random effects can readily be incorporated to cope  
5 with heterogeneity in the detection probability (Huggins, 2001) and deal with a frailty in  
6 the survival probability (Vaupel and Yashin, 1985). Second, the combination of various  
7 sources of information which has recently received a growing interest, (e.g. recovery  
8 and recapture data, Catchpole et al., 1998; recovery and census data, Besbeas et al.,  
9 2002; Besbeas et al., 2003) can now be operated/conducted in a unique SSM framework  
10 and hence benefits from the corpus of related methods. Of particular importance, we  
11 are currently investigating the robust detection of density-dependence by accounting  
12 for error in the measurement of population size using the combination of census data  
13 and data on marked individuals.

14 Because most often, data collected in population dynamics studies give only a noisy  
15 output of the demographic process under investigation, the SSM framework provides  
16 a flexible and integrated framework for fitting a wide range of models which, with  
17 widespread adoption, has the potential to advance significantly ecological statistics  
18 (Buckland et al., 2004; Thomas et al., 2005).

## 1 **5 Acknowledgments**

2 O. Gimenez's research was supported by a Marie-Curie Intra-European Fellowship  
3 within the Sixth European Community Framework Programme. This project was  
4 funded by the Action Incitative Régionale BioSTIC-LR 'Modélisation intégrée en dy-  
5 namique des populations : applications à la gestion et à la conservation'. The authors  
6 would like to warmly thank J.-D. Lebreton for his support during this project. We also  
7 thank a referee whose comments help improving a previous draft of the paper.

## 8 **6 Literature Cited**

- 9 Anderson-Sprecher, R., Ledolter, J., 1991. State-space analysis of wildlife telemetry  
10 data. *Journal of the American Statistical Association*. 86, 596–602.
- 11 Arnason, A.N., 1973. The estimation of population size, migration rates and survival  
12 in a stratified population. *Research in Population Ecology*. 15, 1–8.
- 13 Besbeas, P., Freeman, S.N., Morgan, B.J.T., Catchpole, E.A., 2002. Integrating mark-  
14 recapture-recovery and census data to estimate animal abundance and demographic  
15 parameters. *Biometrics*. 58, 540–547.
- 16 Besbeas, P., Lebreton, J.D., Morgan, B.J.T., 2003. The efficient integration of abun-  
17 dance and demographic data. *Journal of the Royal Statistical Society Series C - Applied*  
18 *Statistics*. 52, 95–102.
- 19 Brooks, S. P., Catchpole, E. A., Morgan, B. J. T., 2000. Bayesian animal survival  
20 estimation. *Statistical Science*. 15, 357–376.
- 21 Brownie, C., Anderson, D.R., Burnham, K.P., Robson, D.S., 1985. *Statistical infer-*  
22 *ence from band recovery data - a handbook*. United States Fish and Wildlife Service,  
23 Washington.

1 Buckland, S.T., Newman, K.B., Thomas, L., Koesters, N.B., 2004. State-space models  
2 for the dynamics of wild animal populations. *Ecological Modelling*. 171, 157–175.

3 Catchpole, E.A., Freeman, S.N., Morgan, B.J.T., Harris, M.P., 1998. Integrated recovery/recapture data analysis. *Biometrics*. 54, 33–46.

5 Choquet, R., Reboulet, A.-M., Pradel, R., Gimenez, O., Lebreton, J.-D., 2004. M-  
6 SURGE: new software specifically designed for multistate capture-recapture models.  
7 *Animal Biodiversity and Conservation*. 27, 207–215.

8 Clark, J.S., 2005. Why environmental scientists are becoming Bayesians. *Ecology*  
9 *Letters*. 8, 2–14.

10 Clark, J.S., Ferraz, G.A., Oguge, N., Hays, H., DiCostanzo, J., 2005. Hierarchical Bayes  
11 for structured, variable populations: From recapture data to life-history prediction.  
12 *Ecology*. 86, 2232–2244.

13 Cormack, R. M., 1964. Estimates of survival from the sighting of marked animals.  
14 *Biometrika*. 51, 429–438.

15 De Valpine, P., 2004. Monte Carlo state-space likelihoods by weighted posterior kernel  
16 density estimation. *Journal of the American Statistical Association*. 99, 523–536.

17 Dupuis, J., 1995. Bayesian estimation of movement and survival probabilities from  
18 capture-recapture data. *Biometrika*. 82, 761–772.

19 Ellison, A.M., 2004. Bayesian inference in ecology. *Ecology Letters*. 7, 509–520.

20 Gimenez, O., Bonner, S., King, R., Parker, R. A., Brooks, S., Jamieson, L. E., Grosbois,  
21 V., Morgan, B. J. T., Thomas, L., submitted. WinBUGS for population ecologists:  
22 Bayesian modeling using Markov Chain Monte Carlo methods. *Environmental and*  
23 *Ecological Statistics*.

24 Gelman, A., 1996. Inference and monitoring convergence. In Gilks, W. R., Richardson,  
25 S. Spiegelhalter, D. J. (eds), *Markov chain Monte Carlo in practice*, pp 131–143.

1 Chapman and Hall, London.

2 Harvey, A.C., 1989. Forecasting, structural time series models and the Kalman Filter.  
3 Cambridge University Press, Cambridge.

4 Huggins, R., 2001. A note on the difficulties associated with the analysis of capture-  
5 recapture experiments with heterogeneous capture probabilities. *Statistics and Proba-*  
6 *bility Letters.* 54, 147–152.

7 Ihaka, R., Gentleman R., 1996. R: a language for data analysis and graphics. *Journal*  
8 *of Computational and Graphical Statistics.* 5, 299–314.

9 Jolly, G. M., 1965. Explicit estimates from capture-recapture data with both death and  
10 immigration-stochastic model. *Biometrika.* 52, 225–247.

11 Lebreton, J.-D., Burnham, K. P., Clobert, J., Anderson, D. R., 1992. Modeling survival  
12 and testing biological hypotheses using marked animals: A unified approach with case  
13 studies. *Ecological Monograph.* 62, 67–118.

14 Lebreton, J.D., Almeras, T., Pradel, R., 1999. Competing events, mixtures of informa-  
15 tion and multistrata recapture models. *Bird Study.* 46, 39–46.

16 Lebreton, J.D., Pradel, R., 2002. Multistate recapture models: modelling incomplete  
17 individual histories. *Journal of Applied Statistics.* 29, 353–369.

18 Millar, R.B., Meyer, R., 2000. Non-linear state space modelling of fisheries biomass  
19 dynamics by using Metropolis-Hastings within-Gibbs sampling. *Journal of the Royal*  
20 *Statistical Society Series C - Applied Statistics.* 49, 327–342.

21 Newman, K. B. 1998. State-space modeling of animal movement and mortality with  
22 application to salmon. *Biometrics.* 54, 1290–1314.

23 Pollock, K.H., 1991. Modeling capture, recapture, and removal statistics for estimation  
24 of demographic parameters for fish and wildlife populations - past, present, and future.  
25 *Journal of the American Statistical Association.* 86, 225–238.

- 1 Pollock, K. H., 2002. The use of auxiliary variables in capture-recapture modelling: an  
2 overview. *Journal of Applied Statistics*. 29, 85–102.
- 3 Pradel, R., 2005. Multievent: An extension of multistate capture-recapture models to  
4 uncertain states. *Biometrics*. 61, 442–447.
- 5 Schwarz, C.J., Schweigert, J.F., Arnason, A.N., 1993. Estimating migration rates using  
6 tag-recovery data. *Biometrics*. 49, 177–193.
- 7 Schwarz, C. J., Seber, G. A. F., 1999. Estimating animal abundance: review III.  
8 *Statistical Science*. 14, 427–56.
- 9 Seber, G. A. F., 1965. A note on the multiple-recapture census. *Biometrika*. 52,  
10 249–259.
- 11 Spiegelhalter, D. J., Thomas, A., Best, N. G., Lunn, D., 2003. WinBUGS user manual  
12 - Version 1.4. MRC Biostatistics Unit, Cambridge, UK.
- 13 Sturtz, S., Ligges, U., Gelman, A., 2005. R2WinBUGS: a package for running Win-  
14 BUGS from R. *Journal of Statistical Software*. 12, 1–16.
- 15 Thomas, L., Buckland, S.T., Newman, K.B., Harwood, J., 2005. A unified framework  
16 for modelling wildlife population dynamics. *Australian and New Zealand Journal of*  
17 *Statistics*. 47, 19–34.
- 18 Vaupel, J.W., Yashin, A.I., 1985. Heterogeneity ruses - some surprising effects of  
19 selection on population dynamics. *American Statistician*. 39, 176–185.
- 20 Wang, G., 2006. On the latent state estimation of nonlinear population dynamics using  
21 Bayesian and non-Bayesian state-space models. *Ecological Modelling*. doi: 10.1016/  
22 j.ecolmodel.2006.09.004.
- 23 Williams, B. K., Nichols, J. D., Conroy, M. J., 2002. Analysis and management of  
24 animal populations. Academic Press, San Diego.



1 **Appendix A: WinBUGS code for fitting the CJS**  
2 **model using the SSM formulation**

```
3 #####  
4 # MODEL #  
5 # State-space formulation of the Cormack-Jolly-Seber model #  
6 # observations = 0 (non-encountered) and 1 (encountered) #  
7 # states = 0 (dead) and 1 (alive) #  
8 #####  
9 model  
10 {  
11 # Define the priors for survival phi and detectability p  
12 p[1] <- 1  
13 phi[1] <- 1  
14 for (j in 2:K)  
15 {  
16 phi[j] ~ dbeta(1,1)  
17 p[j] ~ dbeta(1,1)  
18 }  
19 # Define the SYSTEM PROCESS  
20 for (i in 1:n)  
21 {  
22 # if first capture  
23 PrX[i,e[i]] <- 1 # Pr(alive | first capture) = 1  
24 X[i,e[i]+1] ~ dbern(PrX[i,e[i]]) # alive (a 1 is generated with certainty)
```

```

1 Pr0[i,e[i]] <- X[i,e[i]+1] # detection probability at initial detection is 100%
2 # otherwise
3   for (j in (e[i]+1):K)
4     {
5       PrX[i,j] <- phi[j] * X[i,j]
6       X[i,j+1] ~ dbern(PrX[i,j])
7       Pr0[i,j] <- p[j] * X[i,j+1]
8     }
9 # fullfil the remaining cells with zeros
10  for (j in 1:(e[i]-1))
11    {
12      PrX[i,j] <- 0
13      X[i,j] <- 1
14      Pr0[i,j] <- 0
15    }
16  }
17 # Define the OBSERVATION PROCESS
18 for (h in 1:nx)
19   {
20     data[h,3] ~ dbern(Pr0[data[h,1],data[h,2]])
21   }
22 }
23 #####
24 # DATA
25 # 'K' is the number of encounter occasions

```

```

1 # 'n' is the number of individuals
2 # 'nx' is 'K' times 'n'
3 # 'e' is the vector of first encounters ('n' components)
4 # 'data' is a matrix with dimensions 'nx' times 3 where
5 #   the first column gives the current individual (1,...,'n'),
6 #   the second column gives the current encounter occasion (1,...,'K'),
7 #   the third column gives the observation (= 1 if detection, = 0 otherwise)
8 #   corresponding to the current individual and current encounter occasion
9 #####

```

## 10 Appendix B

11 Let  $Z_{i,t} = [X_{i,t-1}, X_{i,t}]$  be a bivariate random vector where its two components are  
12 denoted  $Z_{i,t}^1$  and  $Z_{i,t}^2$ . Equation (8) becomes

$$13 \quad Y_{i,t}|Z_{i,t} \sim \text{Bernoulli}\left((Z_{i,t}^1 - Z_{i,t}^2)\lambda_{i,t}\right) \quad (10)$$

14 and Equation (9) becomes

$$15 \quad Z_{i,t+1}|Z_{i,t} = \begin{cases} (Z_{i,t+1}^1|Z_{i,t}) = Z_{i,t}^2 \\ Z_{i,t+1}^2|Z_{i,t} \sim \text{Bernoulli}(Z_{i,t}^2\phi_{i,t}). \end{cases} \quad (11)$$

16 The system defined by Equation (10) and Equation (11) is a state-space model and it  
17 is equivalent to the model defined by Equation (8) and Equation (9).

18 Note that an alternative state-space formulation can be adopted using a multistate  
19 formulation of the RR model (Lebreton et al., 1999) and Section 3.1.




Figure 1: Posterior distributions for the survival and detection probabilities using the CJS model applied to the Dipper data set as estimated by the state-space model and MCMC methods.

Table 1: Estimated survival and detection probabilities for the Dipper data using the CJS model and three different methods, the state-space model (SSM) using a Monte Carlo Markov Chain (MCMC) method, the product-multinomial model (PMM) using a MCMC method and the PMM using a maximum-likelihood (ML) method. The two first methods were implemented using program WinBUGS (Spiegelhalter et al., 2003), while program M-SURGE (Choquet et al., 2004) was used to implement the last one.

Parameter	SSM using MCMC	PMM using MCMC	PMM using ML
	Posterior median/mode (SD)	Posterior median/mode (SD)	MLE (SE)
$\phi_1$	0.721/0.722 (0.132)	0.723/0.693 (0.132)	0.718 (0.156)
$\phi_2$	0.448/0.456 (0.071)	0.448/0.460 (0.071)	0.435 (0.069)
$\phi_3$	0.480/0.493 (0.060)	0.480/0.476 (0.061)	0.478 (0.060)
$\phi_4$	0.628/0.624 (0.061)	0.627/0.616 (0.060)	0.626 (0.059)
$\phi_5$	0.602/0.601 (0.057)	0.602/0.607 (0.057)	0.599 (0.056)
$\phi_6$	0.713/0.640 (0.142)	0.720/0.628 (0.143)	- (-)*
$p_2$	0.671/0.658 (0.134)	0.670/0.691 (0.134)	0.696 (0.166)
$p_3$	0.883/0.918 (0.083)	0.883/0.904 (0.083)	0.923 (0.073)
$p_4$	0.888/0.914 (0.063)	0.889/0.912 (0.063)	0.913 (0.058)
$p_5$	0.882/0.885 (0.057)	0.883/0.904 (0.057)	0.901 (0.054)
$p_6$	0.913/0.920 (0.052)	0.912/0.935 (0.051)	0.932 (0.046)
$p_7$	0.735/0.724 (0.142)	0.727/0.648 (0.143)	- (-)*

\* Non-identifiability detected.